

# NIH RELAIS Document Delivery

NIH-10126347

NIH -- W1 BR593

PAMELA GEHRON ROBEY  
CSDB/NIDR/NIH Bldng 30 Rm 228  
30 CONVENT DRIVE MSC 4320  
BETHESDA, MD 20892

ATTN:	SUBMITTED:	2002-02-02 05:11:56
PHONE: 301-496-4563	PRINTED:	2002-02-05 13:37:54
FAX: 301-402-0824	REQUEST NO.:	NIH-10126347
E-MAIL:	SENT VIA:	LOAN DOC 5769534

NIH	Fiche to Paper	Journal
TITLE:	BRITISH JOURNAL OF ORAL MAXILLOFACIAL SURGERY	
PUBLISHER/PLACE:	Churchill Livingstone Edinburgh	
VOLUME/ISSUE/PAGES:	1998 Aug;36(4):264-73	264-73
DATE:	1998	
AUTHOR OF ARTICLE:	Posnick JC	
TITLE OF ARTICLE:	Fibrous dysplasia of the craniomaxillofacial regio	
ISSN:	0266-4356	
OTHER NOS/LETTERS:	Library reports holding volume or year 8405235 9762454	
SOURCE:	PubMed	
CALL NUMBER:	W1 BR593	
REQUESTER INFO:	AB424	
DELIVERY:	E-mail: probey@DIR.NIDCR.NIH.GOV	
REPLY:	Mail:	

NOTICE: THIS MATERIAL MAY BE PROTECTED BY COPYRIGHT LAW (TITLE 17, U.S. CODE)

-----National-Institutes-of-Health,-Bethesda,-MD-----



## Fibrous dysplasia of the craniomaxillofacial region: current clinical perspectives

J. C. Posnick

*Craniomaxillofacial Center, Georgetown Craniofacial Center and Department of Surgery, Pediatrics, Otolaryngology/Head and Neck Surgery and Oral and Maxillofacial Surgery, Georgetown University Medical Center, Washington, DC, USA*

**SUMMARY.** Fibrous dysplasia is a benign fibro-osseous disease of bone of unknown etiology. Its occurrence in the craniomaxillofacial skeleton is frequent and varies in severity from an asymptomatic monostotic lesion to polyostotic involvement resulting in progressive functional deficit and aesthetic problems.

With the advent of refined instrumentation and craniofacial surgical techniques, a more aggressive, non-disabling approach to these benign yet deforming fibro-osseous growths is possible. In some patients, complete excision of the involved bone with graft reconstruction of the resultant defect with primary autogenous bone may be possible. Lifelong continuous ongoing monitoring of the involved region is required throughout the patient's life.

Fibrous dysplasia is a benign fibro-osseous bone disease of unknown aetiology.<sup>1,2</sup> The range of skeletal involvement varies from an asymptomatic monostotic lesion to polyostotic involvement resulting in progressive functional deficit and aesthetic problems.<sup>3-39</sup> As part of the fibro-osseous process normal bone is replaced by cellular fibrous tissue and immature bone.<sup>12,26</sup> The clinical behaviour of fibro-osseous lesions varies greatly, though distinguishing histological features are few.<sup>40</sup>

Key points in the management of a patient with fibrous dysplasia are: the accuracy of the diagnosis; radiographic assessment; clinical evaluation; planning of interventions; procedures carried out; and the long term follow-up. Accurate diagnosis is important to understand better the natural history of the disease and the associated conditions including Albright's syndrome. The radiographic assessment should confirm the diagnosis and allow for assessment of the extent of skeletal involvement. A complete clinical evaluation will record any alteration of breathing, swallowing, speech, occlusion, vision, hearing and cerebral function. Thoughtful planning of the timing and extent of the intervention followed by precise operations when indicated will maximize both function and aesthetics. Lifelong continuing reassessment to monitor changes in the patient's needs throughout the patient's life is required.

### HISTORICAL PERSPECTIVE

Fibrous dysplasia is a congenital dysplastic disease of bone that may occur in a single bone (monostotic) or many bones (polyostotic) and it is considered to

be a type of fibro-osseous lesion.<sup>3,10</sup> Fibrous dysplasia of bone was first described by Von Recklinghausen in 1891 when he incorrectly included two cases of polyostotic fibrous dysplasia with his description of osteitis fibrosa cystica of hyperparathyroidism.<sup>41</sup> In 1937, Albright described a syndrome of polyostotic fibrous dysplasia that included: skeletal changes, cutaneous pigmentation, and endocrine disturbances (precocious puberty in girls being the most striking example).<sup>42</sup> In 1938, Lichtenstein reviewed the publications to date, further delineated the clinical spectrum, and clarified the pathological anatomy of this condition which he named fibrous dysplasia.

### Aetiology and pathophysiology

The cause of the abnormal fibro-osseous process has not been clarified.<sup>43-47</sup> Macroscopically, the lesion looks yellowish or grey and is highly vascular.<sup>26,44</sup> Histopathological analysis shows that it consists of areas of fibrous tissue interwoven with newly formed bone. The fibrous tissue may be extensive or limited and may vary within areas of the same lesion. The bony trabeculae vary in shape. Some are slender and C-shaped (called 'Chinese' characters trabecula). There is no lamellar bone formation in the classic histological picture of the disease.<sup>40</sup>

Why there is maturational arrest at the woven bone stage is open to question.<sup>40</sup> Several authorities have suggested that the lesion is the result of trauma.<sup>43,48</sup> Lichtenstein indicated that it is caused by aberrant activity in the bone-forming mesenchymal tissue.<sup>3</sup> Most authorities consider it to be a non-neoplastic developmental lesion of bone.

### Clinical and radiographic presentation

The process may involve a single bone, more than one bone in a single region (sphenoid wing region) or be disseminated throughout the body,<sup>49</sup> but the monostotic form of the disease is the most common.<sup>5,14</sup> Polyostotic fibrous dysplasia associated with precocious puberty and café-au-lait spots is known as Albright's syndrome.<sup>42,50-51</sup> Although both forms of the disease usually begin in childhood, age of onset and symptoms vary considerably from patient to patient and they usually seek medical attention once they are aware of symptoms or signs. Some are identified only on routine radiographic or physical examination. Harris *et al.* reported a patient with no symptoms whose disease was not diagnosed until he was 68 years of age.<sup>48</sup> At the time of diagnosis, the disease had affected 30% of the skeleton.

In many patients, the lesions are recognized in early childhood, grow slowly, and stabilize in early adult life. The point at which growth is arrested is unpredictable and is not always reached by puberty. Such lesions rarely degenerate into osteosarcomas,<sup>52-57</sup> which are more likely to follow a course of radiotherapy undertaken in an attempt to control the lesion. Radiation is of no value in the treatment of this condition, and the possibility of developing a late post-irradiation sarcoma of bone makes it an absolute contraindication.

The most commonly involved bones of the craniofacial skeleton are the maxilla and frontal bones. When the maxilla is affected, other adjacent bones separated by sutures (zygoma, sphenoid, frontal, nasal) are often also involved and so it is strictly not monostotic. When the frontal bone is affected, involvement of the adjacent sphenoid, temporal, and zygomatic bones is common.

The most usual finding is painless enlargement of the involved bone, which presents as facial asymmetry or a pathological fracture of an extremity.<sup>12,22,29,58-59</sup> The onset is often so gradual that the patient cannot remember when the swelling began. Leads and Seaman described 46 patients with fibrous dysplasia of the skull among whom a painless mass accounted for 45% of the initial symptoms and presentation.<sup>44</sup> Twenty-four percent were detected incidentally by radiography. Symptoms are more common when enlargement of the bones at the base of the skull cause narrowing of the neural foramina leading to blindness, tinnitus and deafness.<sup>23,28,46,60-61</sup> Proliferation of the tumour may also obstruct ostia and produce epiphora in the case of the nasolacrimal apparatus or sinusitis if the nasal ostia are involved. Partial obstruction of the nasal passages may result in altered resonance of speech, diminished nasal airflow with mouth breathing, and sleep apnoea.<sup>59</sup>

Fibrous dysplasia may affect the anterior base of the skull with inferior displacement of the orbital roof which results in proptosis of the eye.<sup>29,32,39,60-63</sup> If the maxilla is primarily involved with extensions to the orbital floor, the globe will be displaced upwards

resulting in orbital dystopia. In either case, the lateral wing of the sphenoid bone and the optic canal may be narrowed, resulting in direct compression of the optic nerve(s) with subsequent papilloedema, optic atrophy, and blindness.

### Differential diagnosis

Cherubism is also a fibro-osseous lesion with several features that distinguish it from fibrous dysplasia of bone. Cherubism is a benign, hereditary (autosomal dominant pattern) giant cell lesion of the jaws that presents in children as a bilateral painless swelling, generally between the ages of 2 and 5 years, with progression of the lesion until puberty when it often regresses spontaneously.<sup>70-74</sup> It generally requires little or no treatment but may displace teeth, distort facial features, and result in pathological fractures. Other developmental disorders of bone often confused with fibrous dysplasia include giant cell granuloma, aneurysmal bone cyst, and florid fibrous dysplasia. Ossifying fibroma is a true bone tumour and should be distinguished from these other lesions.<sup>73-74</sup>

### TREATMENT


Radiotherapy is contraindicated because of its potential for malignant transformation and chemotherapy is not effective, so surgical intervention is the only treatment option. The timing and extent of operation as opposed to conservative observation requires clinical judgement and informed consent.<sup>32-33,39,48,58-59,62-69</sup>

### Case reports


Five patients show the range of presentations and treatments that lend perspective to the clinical management of fibrous dysplasia of the cranio-maxillofacial regions. (Figs 1-5).

### CONCLUSIONS

With the advent of refined instrumentation and craniofacial surgical techniques, a more aggressive, non-disabling approach to those benign yet deforming fibro-osseous growths is possible. In some patients, complete excision of the involved bone with reconstruction of the resultant defect with primary autogenous bone may be possible. For others, despite the unpredictable nature of the residual disease, effective debulking of the regional fibrous dysplasia may be done with improvements in function, facial appearance, and self esteem. Lifelong continuous monitoring is required.




**Fig. 1** A 5-year-old girl with polyostotic fibrous dysplasia and massive involvement of the maxilla and mandible bilaterally. An asymmetric swelling of the mandible (left more than right) was first recognized when the child was 18 months old; it rapidly enlarged and maxillary involvement became evident with deterioration of breathing, chewing, speech and swallowing. Pathological fractures were sustained in all four extremities as a result of involvement in fibrous dysplasia. (A) Preoperative oblique view. (B) Oblique view 10 days after undergoing radical maxillary and mandibular debulking procedure. (C) Frontal three-dimensional craniofacial CT view before operation. (D) Two-dimensional CT view through body and symphyseal region of mandible before operation. (From: Posnick JC, Hughes CA, Milmo G et al: Polyostotic fibrous dysplasia: An unusual presentation in childhood. *J Oral Maxillofac Surg.* Reproduced with permission).




**Fig. 2** – A 3-year-old boy presented to the emergency department with progressive visual loss and a degree of proptosis of the left eye. Both optic nerves were severely atrophic on fundoscopy, particularly in the left proptotic eye. A craniofacial CT and magnetic resonance imaging confirmed the diagnosis of fibrous dysplasia and showed the extent of involvement of the anterior base of the skull and orbits. (A) Frontal view at the time of presentation. (B) Coronal (*left*) and axial (*right*) CT slices through the cranial base and orbits, showing the extent of abnormal fibro-osseous bone resulting in compression of the optic nerve. (C) Close up view of optic chiasm after debulking of tumour. (D) The orbital roofs are reconstructed with autogenous split cranial grafts and fixed in place with microplates and screws to the upper orbital unit before being reinserted. (E) Full-face view 1½ years after operation. (F) Coronal CT close to optic foramen before (*left*) and after (*right*) tumour resection and reconstruction of orbital roofs. (From: Posnick JC, Wells MD, Drake JM et al: Childhood fibrous dysplasia presenting as blindness: A skull base approach for resection and immediate reconstruction. *Ped Neurosurg* 1993; 19: 260–266. Reproduced with permission)






**Fig. 3** – A 17-year-old boy born with an incomplete cleft of the left lip and palate. He underwent orthodontic treatment in preparation for planned orthognathic surgery and bone grafting of his alveolar defect. One month before surgery, he complained of diplopia of the right eye. Radiographs showed fibrous dysplasia of the right maxilla, orbit, zygoma and towards the base of the skull along the orbital floor. He underwent removal of fibrous dysplastic bone of the upper maxilla, anterior zygoma, lateral orbital and infraorbital rim, lateral orbital wall and floor with decompression of the orbital contents. Reconstruction comprised a pedicled temporalis muscle flap used to close the dead space of the maxillary sinus and then full-thickness autogenous cranial bone graft to reconstruct the anterior maxilla, anterior zygoma, lateral and infraorbital rims and lateral orbital wall and floor. (A) Full-face view before operation. (B) Full-face view 1 year after resection and reconstruction. (C) Coronal and axial CT slices through midface and orbit showing extent of fibrous dysplasia.




**Fig. 3 – (cont.)** (D) The orbital contents are decompressed by removal of abnormal fibrous dysplastic bone. (E) Elevation of anterior half of temporalis muscle through coronal incision. (F) View of temporalis muscle flap obliterating dead space of maxillary sinus with overlying reconstruction of skeletal defect with cranial bone.




**Fig. 4** – An 8-year-old girl had a left fronto-orbital and sphenoid wing monostotic fibrous dysplasia which presented as a painless, hard mass, followed by diplopia and headache. An ophthalmological assessment and CT scan confirmed the extent of involvement. (A) Close-up facial view showing left frontotemporal mass. (B) Bone scan confirms the area of reactivity of the left fronto-orbital and sphenoid wing region. (C) View through the coronal incision showing the mass in the left fronto-orbital and sphenoid wing mass.






**Fig. 4 - (cont.)** (D) The tumour mass is resected back to the optic foramen. (E) Cranial bone graft reconstruction. Stabilization is achieved with multiple titanium bone plates and screws. (F) Frontal view shows the patient 4 years after resection and reconstruction. (G) Two-dimensional CT scan views through orbit demonstrate reconstructed region. (From: Posnick JC: The role of plate and screw fixation in the management of pediatric head and neck tumours. In: Gruss JS, Manson PM, Yaremchuk MJ, eds. Rigid fixation of the craniomaxillofacial skeleton. Stoneham, MA, Butterworths, 1992: 665. Reproduced with permission).



**Fig. 5** – A 6-year-old child presented with a firm mass over the upper forehead. At 2½ years of age, he underwent removal of a bony mass from the same region with immediate rib graft reconstruction by another surgeon. After assessment including craniofacial, neurosurgical, ophthalmological evaluation and CT scan, recurrence of the fibrous dysplasia of bone was removed and immediately reconstructed with autogenous cranial bone. (A) Preoperative profile. (B) Profile 2 months after resection and immediate reconstruction. (C) Intraoperative view of cranial vault showing the mass.



**Fig. 5 – (cont.)** (D) Intraoperative view after removal of the left frontotemporal mass and harvesting of full-thickness right occipitoparietal cranial bone for reconstruction. (E) View of cranial vault with donor (frontotemporal) and recipient (occipitoparietal) sites reconstructed with split autogenous cranial grafts fixed with microplates and screws.

## References

- Gorlin RJ, Cohen MM, Levin LS (eds). *Syndromes of the head and neck*, 3rd ed., New York: Oxford University Press, 1990.
- Eversole LR, Sabes WR, Rovin S. Fibrous dysplasia: a nosologic problem in the diagnosis of fibro-osseous lesions of the jaws. *J Oral Pathol Med* 1972; 1: 189–220.
- Lichtenstein L. Polyostotic fibrous dysplasia. *Arch Surg* 1938; 36: 874–898.
- Lichtenstein L, Jaffe HL. Fibrous dysplasia of bone. *Arch Pathol* 1942; 33: 777–816.
- Schlumberger HG. Monostotic fibrous dysplasia. *Mil Surg* 1946; 99: 504–527.
- Schlumberger HG. Fibrous dysplasia (ossifying fibroma) of the maxilla and mandible. *Am J Ortho Oral Surg (Oral Surg Sec)* 1946; 32: 579–587.
- Windolz F. Cranial manifestations of fibrous dysplasia of bone. *Am J Roentgenol Rad Ther* 1947; 58: 51–63.
- Fisher AI. Fibrous dysplasia of facial bones. *J Oral Surg* 1948; 6: 218–227.
- Feiring W, Feiring EH, Davidoff LM. Fibrous dysplasia of the skull, 1950.
- Kaufman M. Fibrous dysplasia of bone; monostotic and polyostotic forms. *J Oral Surg* 1951; 9: 93–104.
- Weyland RD, Craig WM, Rucker CW. Unusual lesions involving the optic chiasm. *Proc Staff Meet Mayo Clin* 1952; 27: 505–511.
- Thoma KH. Differential diagnosis of fibrous dysplasia and fibro-osseous neoplastic lesions of the jaws and their treatment. *J Oral Surg* 1956; 14: 185–194.
- Davis MI, Yurdley JH. Fibrous dysplasia of bone. *Am J Med Sci* 1957; 234: 590–606.
- Cooke BED. Benign fibro-osseous enlargement of the jaws. *Br Dent J* 1957; 102: 1–14.
- Church LE. Polyostotic fibrous dysplasia of bone. *Oral Surg* 1958; 11: 184–196.
- Sherman RS, Glauser OJ. Radiologic identification of fibrous dysplasia of the jaws. *Radiology* 1958; 71: 553–558.
- Zimmerman DC, Dahlin DC, Stafne EC. Fibrous dysplasia of the maxilla and mandible. *Oral Surg* 1958; 11: 55–68.
- Lucas RB. Fibrous dysplasia in the jaws. *J R Coll Surg Edinb* 1962; 7: 255–267.
- Reed RJ. Fibrous dysplasia of bone. A review of 25 cases. *Arch Pathol* 1963; 75: 480–495.
- Van Horne PE. Fibrous dysplasia: a clinical pathologic study of orthopedic surgery cases. *Mayo Clin Proc* 1963; 38: 175–189.
- Houston WO Jr. Fibrous dysplasia of maxilla and mandible; clinico-pathologic study and comparison of facial bone lesions with lesions affecting general skeleton. *J Oral Surg* 1965; 23: 17–39.
- Ramsey HE, Strong EW, Faxell EL. Fibrous dysplasia of the craniofacial bones. *Am J Surg* 1968; 116: 542–547.
- Sassin JF, Rosenberg RN. Neurological complications of fibrous dysplasia of the skull. *Arch Neurol* 1968; 18: 363–369.
- Frut D, Stutzman L. Fibrous dysplasia of bone, review of twenty-four cases. *Am J Med* 1968; 44: 421–429.
- Dahlgren SE, Lind PO, Lindbom A et al. Fibrous dysplasia of jaw bones. A clinical, roentgenographic and histopathologic study. *Acta Otolaryngol* 1969; 68: 257–270.
- Waldron CA, Giansanti JS. Benign fibro-osseous lesions of the jaws: A clinical-radiologic-histologic review of sixty-five cases. 1. Fibrous dysplasia of the jaws. *Oral Surg Oral Med Oral Pathol* 1973; 35: 190–201.

27. Obwegeser HL, Hans P, Freihofer M et al. Variations of fibrous dysplasia in the jaws. *J Maxillofac Surg* 1973; 1: 161-171.
28. Finney HL, Roberts TS. Fibrous dysplasia of the skull with progressive cranial nerve involvement. *Surg Neurol* 1976; 6: 341-343.
29. Schonder A. Fibrous dysplasia of bone with proptosis. *Am J Dis Child* 1977; 131: 678-679.
30. El Deeb M, Waite DE, Gorlin RJ. Congenital monostotic fibrous dysplasia - a new possibly autosomal recessive disorder. *J Oral Surg* 1979; 37: 520-525.
31. Adekee EO, Edwards MB, Goubran GF. Fibro-osseous lesions of the skull, face and jaws in Kaduna, Nigeria. *Br J Oral Surg* 1980; 18: 57-72.
32. Ajagbe HA, Daramola JO. Fibro-osseous lesions of the jaw. *Journal of National Medical Association*, 1983; 75: 593-598.
33. Moore AT, Buncie JR, Munro IR. Fibrous dysplasia of the orbit in childhood. Clinical features and management. *Ophthalmology* 1985; 92: 12-20.
34. Chen Y-R, Fairholm D. Fronto-orbito-sphenoid fibrous dysplasia. *Ann Plast Surg* 1985; 15: 190-202.
35. Waldron CA. Fibro-osseous lesions of the jaws. *J Oral Maxillofac Surg* 1985; 43: 249-262.
36. Lello GE, Sparrow OC. Craniofacial polyostotic fibrous dysplasia. *J Maxillofac Surg* 1985; 13: 267-272.
37. Kreutziger KL. Giant fibrous dysplasia of ten mandibles: Surgical management. *Laryngoscope* 1989; 99: 618-630.
38. Hara H, Ohishi M, Higuchi Y. Fibrous dysplasia of the mandible associated with large solitary bone cyst. *J Oral Maxillofac Surg* 1990; 48: 88-91.
39. Posnick JC, Wells MD, Buncie Jr et al. Childhood fibrous dysplasia presenting as blindness: A skull base approach for resection and immediate reconstruction. *Pediatr Neurosurg* 1993; 19: 260-266.
40. Reed RJ. Observations of bone maturation and fibrous dysplasia of bone. *Bull Tulane Med Fac* 1959; 19: 45-53.
41. von Recklinghausen F. Die fibrose oder deformierende osteite. *Festschrift R. Virchow zu seinem 71 Geburtstag*. Berlin, 1891.
42. Albright F, Butler AM, Hampton AO, Smith P. Syndrome characterized by osteitis fibrosa disseminata, areas of pigmentation and endocrine dysfunction, with precocious puberty in females; report of 5 cases. *N Engl J Med* 1937; 216: 727-746.
43. Robinson HBG. Osseous dysplasia: Resection of bone to injury. *J Oral Surg* 1956; 14: 3-14.
44. Leeds N, Seaman WB. Fibrous dysplasia of the skull and its differential diagnosis. A clinical and roentgenographic study of 46 cases. *Radiology* 1962; 78: 570-585.
45. Albright A. Polyostotic fibrous dysplasia; A defence of the entity. *J Clin Endocrinol* 1962; 7: 307-324.
46. Vantilburg W. Fibrous dysplasia. *Handbook of Clin Neurology*, Vol 14. Amsterdam: Elsevier.
47. Gorlin RJ, Chaudhry AP. Oral melanotic pigmentation in polyostotic fibrous dysplasia - Albright's syndrome. *Oral Surg* 1957; 10: 857-862.
48. Harris WH, Dudley HR, Barry RJ. The natural history of fibrous dysplasia. *J Bone Joint Surg* 1962; 44A: 207-233.
49. Fries JW. Roentgen features of fibrous dysplasia of the skull and facial bones. A critical analysis of thirty-nine pathologically proved cases. *Am J Roent* 1957; 77: 71-88.
50. Falconer MA, Cope CI, Robb-Smith AHT. Fibrous dysplasia of bone with endocrine disorders and cutaneous pigmentation (Albright's Disease). *Q J Med* 11: 121-154.
51. Benedict PA. Sex precocity and polyostotic fibrous dysplasia. *Am J Dis Child* 1966; 111: 426-429.
52. Tanner HC, Dahlin DC, Childs DS. Sarcoma complicating fibrous dysplasia. Probable role of radiation therapy. *Oral Surg Oral Med Oral Pathol* 1961; 14: 837-846.
53. Schwartz DT, Alpert M. The malignant transformation of fibrous dysplasia. *Am J Med Sci* 1964; 1: 35/1.
54. Gross CW, Montgomery W. Fibrous dysplasia and malignant degeneration. *Arch Otolaryngol* 1967; 85: 653-657.
55. Slow IA, Stern D, Friedman E. Osteogenic sarcoma arising in a pre-existing fibrous dysplasia: Report of a case. *J Oral Surg* 1971; 29: 126-129.
56. Schwartz DT, Alpert M. The malignant transformation of fibrous dysplasia. *J Bone Joint Surg* 1972; 54-A: 1047-1056.
57. Huvois AG, Higinbotham NL, Miller TR. Bone sarcoma arising in fibrous dysplasia. *J Bone Joint Surg* 1972; 54B: 1047-1056.
58. Ragab MA, Mathog RH. Surgery of massive fibrous dysplasia and osteoma of the midface. *Head Neck Surg* 1987; 9: 202-210.
59. Stompro BE, Bunkis J. Surgical treatment of nasal obstruction secondary to craniofacial fibrous dysplasia. *Plast Reconstr Surg* 1990; 85: 107-111.
60. Gyldensted C, Lester J, Fledelius H. Computed tomography of orbital lesions: A radiological study of 144 cases. *Neuroradiology* 1977; 13: 141-50.
61. Henderson JW. *Orbital Tumors*. 2nd ed. New York: Decker, 1980: 70.
62. Munro IR, Chen Y-R. Radical treatment for fronto-orbital fibrous dysplasia: The chain link principle. *Plast Reconstr Surg* 1981; 67: 719-729.
63. Jackson IT, Armstrong HH, Gomuwka PK et al. Treatment of cranio-orbital fibrous dysplasia. *J Maxillofac Surg* 1982; 10: 138-141.
64. Derome PJ. The transbasal approach to tumors invading the base of the skull. In: Schmidek HH, Sweet WH (eds). *Current Techniques in Operative Neurosurgery*. New York: Grune & Stratton, 1977: 223.
65. Chen Y-R, Noordhoff MS. Treatment of craniomaxillofacial fibrous dysplasia: How early and how extensive? *Plast Reconstr Surg* 1990; 86: 835-842.
66. Munro IR. Discussion: Treatment of craniomaxillofacial fibrous dysplasia: How early and how extensive? *Plast Reconstr Surg* 1990; 86: 843-844.
67. Posnick JC. The role of plate and screw fixation in the treatment of craniofacial malformations. In: *Rigid Fixation of the Craniomaxillofacial Skeleton*, Gruss JS, Manson PM, Yaremchuk MJ (eds). Stoneham, Massachusetts: Butterworth, 1992: 40: 512-526.
68. DePalma AF, Smythe VL. Recurrent fibrous dysplasia in a cortical bone graft. *Clin Orthop* 1963; 26: 136-144.
69. Middleton DS. The place of surgery in fibrous dysplasia of the jaws. *J R Coll Surg Edinb* 1963; 8: 310-318.
70. Reade PC, McKellar GM, Raddan BG. Unilateral mandibular cherubism: Brief review and case report. *Br J Oral Maxillofac Surg* 1984; 22: 189-194.
71. Zachariades N, Papanicolaou S, Xypolyta A et al. Cherubism. *Int J Oral Surg* 1985; 14: 138-145.
72. Zohar Y, Grausbord R, Shabtai F et al. Fibrous dysplasia and cherubism as an hereditary familial disease. *J Craniomaxillofac Surg* 1989; 17: 340-344.
73. Slootweg P, Muller H. Differential diagnosis of fibro-osseous jaw lesions. *J Craniomaxillofac Surg* 1990; 18: 210-214.
74. Reed RJ, Hagy DM. Benign nonodontogenic fibro-osseous lesions of the skull: report of two cases. *Oral Surg* 1965; 19: 214-227.

## The Author

**Jeffrey C. Posnick DMD, MD**

Chief  
Craniomaxillofacial Center  
Director  
Georgetown Craniofacial Center  
Professor  
Department of Surgery, Pediatrics, Otolaryngology/Head and Neck Surgery and Oral and Maxillofacial Surgery  
Georgetown University Medical Center  
Washington DC, USA

Correspondence to Professor J. C. Posnick

Paper received 30 April 1996

Accepted 16 March 1997